

33. We thank T. Sauer, R. Guimerà, M. Sales-Pardo, M. J. Stringer, E. N. Sawardecker, J. Duch, and P. McMullen for insightful comments and suggestions. D.B.S. acknowledges the support of a Consejo Superior de Investigaciones Científicas–Junta para la Ampliación de Estudios (JAE) postdoctoral fellowship. A.S.L.O.C. acknowledges the support of a Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (Brazil)

doctoral fellowship. L.A.N.A. gratefully acknowledges the support of NSF award SBE 0624318. All figures were generated with PyGrace (<http://pygrace.sourceforge.net>) with color schemes from <http://colorbrewer.org>.

Supporting Online Material

www.sciencemag.org/cgi/content/full/325/5948/1696/DC1
Materials and Methods

SOM Text
Figs. S1 to S5
Tables S1 to S3
References

6 April 2009; accepted 5 August 2009
10.1126/science.1174562

Antennal Circadian Clocks Coordinate Sun Compass Orientation in Migratory Monarch Butterflies

Christine Merlin, Robert J. Gegear, Steven M. Reppert*

During their fall migration, Eastern North American monarch butterflies (*Danaus plexippus*) use a time-compensated Sun compass to aid navigation to their overwintering grounds in central Mexico. It has been assumed that the circadian clock that provides time compensation resides in the brain, although this assumption has never been examined directly. Here, we show that the antennae are necessary for proper time-compensated Sun compass orientation in migratory monarch butterflies, that antennal clocks exist in monarchs, and that they likely provide the primary timing mechanism for Sun compass orientation. These unexpected findings pose a novel function for the antennae and open a new line of investigation into clock-compass connections that may extend widely to other insects that use this orientation mechanism.

Eastern North American monarch butterflies, *Danaus plexippus*, undergo one of the most magnificent long-distance mi-

grations observed in animals. Each fall in the northern United States and southern Canada, migratory monarchs travel distances up to 4000 km

to arrive at their overwintering grounds in central Mexico (1, 2). The navigational abilities of the migrants include the use of a time-compensated Sun compass (3–5). Previous studies show that a circadian clock provides the internal timing device that allows the butterflies to correct their flight orientation, relative to skylight parameters, and to maintain a southerly flight bearing as the Sun moves across the sky during the day (3–5).

A distinctive circadian clock mechanism has been recently elucidated in the monarch butterfly (6). It relies on a negative transcriptional feedback loop that involves the transcription factors CLOCK (CLK) and CYCLE (CYC), which drive the expression of *period* (*per*), *timeless* (*tim*), and a vertebrate-like *cryptochrome* designated *cry2*. The translated PER, TIM, and CRY2 proteins

Department of Neurobiology, University of Massachusetts Medical School, Worcester, MA 01605, USA.

*To whom correspondence should be addressed. E-mail: Steven.Reppert@umassmed.edu




Fig. 1. Antennae are necessary for time-compensated sun compass orientation. **(A)** Flight orientation of intact migrants under different lighting conditions. Butterflies were flown between 1100 and 1500 hours from 24 September to 18 October 2008. The large circle represents the 360° of possible directions (0° is north); small solid circles on the perimeter represent the flight orientation of individual butterflies. The arrow indicates the mean vector; arrow length, *r* value. Left, orientation data of butterflies in LD. Right, orientation

data of butterflies in 6-hour-delayed LD. **(B)** Orientation of antennaeless migrants under the different lighting conditions. **(C)** Free-flight behavior of intact migrants (left bar) and those without antennae (right bar). **(D)** Temporal profiles of *per* and *tim* mRNA levels in brains of monarchs with antennae (blue) and without antennae (red). Values are mean ± SEM of three brains. Points at CT0 are replotted at CT24 to show 24-hour trend. Horizontal bars: open, light; black, darkness. *P* values, one-way analysis of variance (ANOVA).

form complexes in the cytoplasm and, after the appropriate time delay, translocate back into the nucleus where CRY2 represses CLK:CYC-mediated transcription (6–8). A *Drosophila*-like CRY also exists in the butterfly, designated CRY1, which functions as a blue light photoreceptor to synchronize (entrain) the circadian clockwork to the prevailing light-dark conditions (6). Four cells in the dorsolateral region of the central brain (the pars lateralis) house the major circadian clocks in butterfly brains (6, 9).

Because circadian rhythms in locomotor activity and the timing of adult eclosion in insects such as silkworms (10) and *Drosophila* (11) are under the control of brain clocks, it has been assumed that the clock involved in time-compensated sun compass orientation in monarchs is also located in the brain (6, 9, 12). This assumption has never been examined directly. The location of the sun compass, on the other hand, is more firmly established. On the basis of electrophysiological studies

in locusts and crickets (13, 14) and genetic studies in *Drosophila* (15), it resides in the central complex, a midline structure in the brain of insects.

In addition to the brain, circadian clocks are also found in the antennae of insects (16–19) and likely exist in the antennae of monarch butterflies. Insect antennal clocks are believed to modulate olfactory reception within the antennae themselves (20, 21), and there has been scant evidence that antennal clocks directly influence brain mechanisms. However, Urquhart presented anecdotal evidence almost 50 years ago suggesting a role of the antennae in the flight orientation of migratory monarchs (2) that was not pursued. In view of this historical observation and our interest in Sun compass mechanisms (12), we rigorously examined the role of the antennae in Sun compass orientation.


To begin, we compared time-compensated Sun compass orientation of intact migrants with migrants whose antennal flagellum had been surgically removed (fig. S1) (22). Both intact and

antennaeless migratory monarchs were housed indoors in either a 12-hour light:12-hour dark (LD) cycle that was timed to coincide with prevailing lighting conditions or a 6-hour-delayed LD cycle. Six to 8 days later, the butterflies housed in either lighting cycle were tethered, and over the next 26 days individual flight direction and group orientation were examined in butterflies flown outdoors in a flight simulator (22, 23) (fig. S2).

Intact monarch migrants housed under LD conditions exhibited directional flight that was oriented as a group significantly to the southwest with a mean vector (α) of 211° ($n = 10$, $r = 0.67$, $P < 0.01$; Rayleigh test; Fig. 1A left), in close agreement with previous reports (3, 4, 24). The group of intact migrants housed under the 6-hour-delayed LD cycle were also oriented significantly but in the northwesterly direction, with an α of 355° ($n = 13$, $r = 0.63$, $P < 0.005$; Fig. 1A, right). The clockwise shift in the direction of orientation in the 6-hour-delayed LD group, relative to the LD

Fig. 2. Circadian clocks in monarch antennae.

(A) Clock gene expression in monarch tissues. Tissues were collected at ZT6 (white) and ZT18 (black). Values are normalized to those in the brain at ZT18 and are mean \pm SEM of four animals. P values, Student's t test: *** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$. **(B)** Clock gene mRNA profiles in antennae. Values are relative to the minimal level for each gene and are the mean \pm SEM of four antennae. Points at CT0 are replotted at CT24. Horizontal bars: gray, subjective day; black, subjective night. P values, one-way ANOVA. **(C)** Circadian profiles of clock protein abundance in antennae. Top, representative autoradiographs in DD. Arrowhead, CRY2 band; the lower band is nonspecific, as shown previously (6). Bottom four graphs, quantification of relative protein levels. Values are normalized to the minimal level of protein expression and are mean \pm SEM of three or four antennae. P values, one-way ANOVA. **(D)** Light-sensitivity of antennae in culture. Top, experimental scheme in LD or in phase-reversed LD (R-LD). Arrows, collection times. Bottom, Western blot analyses. Left, representative blots; right, quantifications. Open bars, mid-light; black bars, mid-dark. Values are mean \pm SEM of five anten-



nae. P values, Student's t test: ** $P < 0.01$ and * $P < 0.05$. **(E)** Daily and circadian patterns of TIM abundance in two sets of cultured antennae (gray and black lines, respectively). Top, lighting conditions. Bottom, representative Western blot.

group, was expected for a time-compensated Sun compass that has been delayed by several hours. However, the magnitude of the difference between the two groups (a clockwise shift of 144° ; $F_{1,21} = 31.92$, $P < 0.0001$; Watson-Williams test) was greater than expected for the 6-hour shift ($\leq 132^\circ$; the speed of the Sun's azimuth varied from 14° to 22° per hour during the study period) but within the range of directions found in a large number of phase-delayed migrants (24).

Remarkably, group flight was disoriented in the antennaeless monarchs studied under either lighting cycle (Fig. 1B). Individual antennaeless migrants housed under either LD or the 6-hour-delayed LD cycle each exhibited significant directional flight. However, orientation of each of the two groups was randomly distributed over the 360° of direction ($n = 13$, $r = 0.23$, $P > 0.05$ for the LD group and $n = 15$, $r = 0.209$, $P > 0.05$ for the 6-hour-delayed LD group) (Fig. 1B). Antennaeless migrants flew as strongly and consistently as intact butterflies in the flight simulator, and the proportion of antennaeless migrants eliminated from analysis because of nondirectional flight did not differ from the proportion of intact migrants (22).

Because the antennae have been shown to mediate flight stability in moths through mechanosensors located at the base of the antennae (25), we also analyzed the effect of antennal flagellum amputation on the free-flight performance of migrant monarchs (22). We found that the majority of intact and antennaeless migrants exhibited normal upward flight (8/10 and 9/10, respectively) (Fig. 1C). Thus, the effect of antennal amputation on flight orientation of migrating monarch butterflies was not the consequence of disrupted flight stability. Taken together, our results show that the antennae are necessary for time-compensated Sun compass orientation in migratory monarch butterflies.

The disorientation of flight behavior in the antennaeless butterflies suggested that the timing component of the time-compensated Sun compass mechanism was disrupted when antennal input was lacking. Is it possible that the antennae are necessary for the proper workings of the molecular clocks in the brain? To test this, we examined the temporal patterns of clock gene expression in the brains of butterflies with and without antennae. With use of the quantitative real-time polymerase chain reaction (QPCR), we analyzed the temporal patterns of the clock genes *per* and *tim* (22); these genes were chosen because they exhibit the most robust mRNA cycling among the clock genes previously studied in monarch heads (6). The butterflies were studied in LD to match the condition used for the flight orientation experiments.

The levels of *per* and *tim* mRNA in brains of intact and antennaeless butterflies cycled in phase (Fig. 1D), as previously described in monarch heads and in DpN1 cells, a monarch butterfly cell line that contains a light-driven clock (6). Thus, compared to the brains from butterflies with in-

tact antennae, the relative timing of *per* and *tim* mRNA levels to the LD cycle were unaltered in antennaeless butterflies. These results suggest that the brain clocks are functioning in synchrony in LD without attached antennae and raise the interesting prospect that the clock for time compensation may actually reside in the antennae and not in the brain.

Accordingly, we next surveyed rhythmic clock gene mRNA and protein expression in the antennae, as well as in other peripheral tissues (legs, thorax, and abdomen) of monarch butterflies. By using QPCR, we found that the mRNA amounts of *per*, *tim*, and *cry2* were significantly lower at mid-light [Zeitgeber time (ZT)6] than at mid-dark (ZT18) in the antennae and the legs, similar to those in the brain (Fig. 2A). The expression of *tim* was also significantly reduced at ZT6 compared with ZT18 in the thorax and the

abdomen, and *per* expression showed a similar oscillation in the abdomen. There were no significant differences in *cry1* mRNA amounts in most tissues examined (Fig. 2A). At the protein level (22), the antenna was the only peripheral tissue of those examined to express clock protein abundance patterns similar to those found in the brain (fig. S3).

Focusing on antennae, a more detailed temporal analysis showed that *per*, *tim*, and *cry2* mRNA levels exhibited robust circadian rhythms in constant darkness (DD) (Fig. 2B), similar to those previously described in monarch brains (Fig. 1D) and/or in DpN1 cells (6). PER and TIM also exhibited significant circadian oscillations in abundance (Fig. 2C). In addition, PER showed temporal changes in electrophoretic mobility corresponding to changes in phosphorylation (6) (Fig. 2C). CRY2 showed a temporal trend in abundance




Fig. 3. Blinding antennal clocks alters their timing. (A) Irradiance curves for different painting conditions. Light measurements were taken under full-spectrum light through plastic that was either painted or not. (B) Light sensitivity of TIM abundance. Top, experimental paradigm. Painted antennae were harvested at CT18 (arrows). Middle, blot of TIM levels from pooled antennae painted clear (C) or black (B) from butterflies in dark or light pulsed. Bottom, quantifications. Values are mean ± SEM of three or four antennae. White bars, clear-painted antennae; black bars, black-painted antennae. P values, Student's t test: ** $P < 0.005$. (C) Temporal patterns of *per* and *tim* expression in brain (left column) and antenna (right column) from butterflies with the antennae painted clear or black. Values are mean ± SEM of three animals, except for the three points without error bars that represent the mean of two animals. Box shading: open, light; dark gray, night; and light gray, subjective day. P values, one-way ANOVA.

that was not significant (Fig. 2C). Circadian cycling of mRNA and protein levels of these core clock components *in vivo* suggests the presence of circadian clocks in the monarch butterfly antenna.

To show that the monarch antennae actually house light-entrained and tissue-autonomous circadian clocks, we examined whether the antennal clocks are reset by light and continue to oscillate when explanted *in vitro* (22). The light sensitivity of isolated antennae was evaluated by maintaining antennae in culture in two oppositely phased LD cycles and probing TIM abundance by Western blot. In both lighting conditions, TIM expression was significantly lower during the light phase (at ZT6) than in the dark (at ZT18) (Fig. 2D). These data show that the LD oscillation in TIM abundance persists *in vitro* in a phase-appropriate manner, suggesting that the antennal clocks can be directly entrained by light, even when disconnected from the brain. We also investigated the ability of the antennae to maintain self-sustained circadian oscillations by analyzing the temporal abundance of TIM in LD and during the first day in DD from individual antennae maintained in culture. TIM abundances oscillated in LD and continued to oscillate, although with reduced amplitude, on the first day in DD (Fig. 2E). Thus, monarch antennae possess light-sensitive circadian clocks, which could function independently from the brain as time-compensation components for sun compass orientation.

If antennal clocks are involved in Sun compass orientation in migrants, then altering their rhythmicity *in vivo* should alter time-compensated sun compass orientation. Blocking light input to antennal clocks should alter their rhythmicity in two ways. First, antennal clocks would continue to oscillate but would gradually drift out of their normal phase relationship with the prevailing lighting cycle (i.e., “free-running” clocks). Second, after several days without light input, the individual free-running clocks would eventually desynchronize from each other because of the innate difference in free-running period length (26).

To prevent light input to intact antennae, we painted the flagellum with an enamel-based black paint (fig. S1) (22), which blocked antennal perception of full-spectrum light (from 300 nm to 800 nm; Fig. 3A); the control, enamel-based clear paint, did not reduce either the intensity or wavelengths of light that could pass through the antenna (Fig. 3A). We verified the efficiency of black paint to block light input *in vivo* by examining the light-induced decline in TIM abundance in painted antennae. As expected, clear- and black-painted antennae in DD had similar TIM abundance during mid-subjective night (the period the lights normally would have been off in LD) (Fig. 3B). However, when both groups were exposed to a 2-hour light pulse from [circadian time (CT)16] to CT18, TIM abundance was substantially lower in the clear-painted antennae

compared with that of antennae painted black (Fig. 3B). Thus, black paint blocks light input to the antennal clocks.

Light sensitivity of the antennae is unlikely to be mediated by opsins because QPCR of the three opsin genes in monarchs [ultraviolet, blue, and long wavelength (9)] showed that they were not detected in the antennae (fig. S4). CRY1 is likely the photoreceptor for the light-input pathway to the antennal clocks and for causing TIM degradation (6). In contrast to *Drosophila* CRY, monarch CRY1 is not degraded by light in either the brain (6) or the antennae (fig. S5). However, knocking down CRY1 expression by RNA interference in DpN1 cells blocks the ability of light to degrade TIM, showing that the monarch CRY1 is a light sensor (6).

To examine how the rhythmicity of antennal clocks is altered by black paint, we quantified by QPCR the expression of the clock genes *per* and *tim* from both clear- and black-painted antennae of butterflies maintained in LD for 11 days after painting (Fig. 3C). In clear-coated antennae, *per* and *tim* mRNA amounts exhibited robust daily rhythms that were in phase with each other. In the same animals, *per* was also rhythmically expressed in the brain, with the phase of the oscillation similar to that found in the antennae. The daily pattern of *tim* in the brain, however, was similar but not significantly rhythmic. The lack of a significant oscillation for *tim* was likely due to the combined effects of individual variation and the shallow





Fig. 4. Blinding antennal clocks alters Sun compass orientation. (A) Flight orientation of migrant butterflies with intact antennae (control, top) and with antennae painted clear (middle) or black (bottom). Butterflies were flown between 1100 and 1500 hours from 20 October to 16 November 2008. Left, butterflies housed in LD. Right, butterflies housed in 6-hour-delayed LD. (B) Relationship

between orientation angle and day of study. Individual orientation directions were standardized to the mean vector of control butterflies ($144^\circ = 360^\circ$) and assumed to be drifting from the mean in a counterclockwise direction over time. Thus, orientation angles are expected to decrease with increasing days after painting, from 360° to 0° to -360° for two revolutions. One orientation direction on day 16 was calculated to be -82° [for the 62° value in (A), lower right]. The absolute value of 82° was used so that all orientations could be tested from 360° to 0° . Black dots, LD 0600 to 1800; open dots, 1200 to 2400. *P* value, linear regression analysis.

amplitude rhythms normally found in the brain (Fig. 1D right) (6). Predictably, *per* and *tim* mRNA amounts were not significantly rhythmic in the black-painted antennae and exhibited peak amounts occurring earlier and for a broader duration than in the clear-painted antennae (Fig. 3C), whereas these genes cycled normally in the brains of the same butterflies.

Painting the antennae black thus appears to specifically block LD entrainment of the antennal clocks. The lack of overall rhythmicity observed in black-painted antennae is likely due to desynchrony among the free-running clocks after 11 days in virtual DD. This apparent arrhythmicity could be due to desynchrony of antennal clocks within the antenna of individual butterflies, between the two antennae, among the antennae of the different butterflies, or a combination of these effects. Thus, sufficient synchrony among free-running clock gene oscillations may exist after several days to synchronize group flight orientation (see below).

To test a potential role of antennal clocks in time-compensated Sun compass orientation of migratory monarchs, we analyzed the flight orientation of migrants with either clear- or black-painted antennae housed in LD or placed from LD into the 6-hour phase-delayed LD cycle after painting. We found that the control migrants without antennal painting and migrants with clear-painted antennae both oriented significantly to the south to southeast with α values of 144° and 162° , respectively ($n = 11$, $r = 0.80$, $P < 0.0001$ and $n = 13$, $r = 0.66$, $P < 0.005$; Fig. 4A). The mean flight orientation did not differ significantly between groups ($F_{1, 22} = 0.77$, $P > 0.05$). Such a south/southeasterly orientation direction has been reported previously for flight simulator studies (27) and free-flight studies late in the season (28). The migrants in our study oriented in the southwesterly direction at the beginning of the fall (Fig. 1 top left) but oriented to the southeast later in the season (Fig. 4A left). Nonetheless, migrants with clear-painted antennae housed under 6-hour-shifted LD showed the appropriate shift in their orientation toward the west ($\alpha = 269^\circ$, $n = 9$, $r = 0.63$, $P < 0.05$; Fig. 4A right); the direction and magnitude of the group orientation differences between the two groups with clear-painted antennae (a clockwise shift of 107° , $F_{1, 20} = 17.48$; $P < 0.001$) were those expected for a time-compensated Sun compass delayed by 6 hours. Thus, although clear-painted antennae lack olfactory reception (fig. S6), they have normal light reception for entraining antennal clocks, and, correspondingly, those migrants show proper time-compensated Sun compass orientation.

A completely different situation was found for flight orientation in the migrants with black-painted antennae. As a group, those migrants housed under LD oriented significantly to the north to northwest ($\alpha = 344^\circ$, $n = 12$, $r = 0.73$, $P < 0.001$; Fig. 4A left), an orientation direction that was 182° different from that of the group of migrants with clear-painted antennae

($F_{1, 23} = 61.17$, $P < 0.00001$). Migrants with black-painted antennae and housed under the 6-hour-shifted LD cycle did not exhibit significant group orientation ($n = 9$, $r = 0.5$, $P > 0.05$; Fig. 4A right), but there was a trend to orient to the southwest. A second difference between the group maintained in LD and the 6-hour-shifted LD group was related to the interval from antennal painting to analysis in the flight simulator. Although the butterflies were flown randomly, retrospectively, the groups differed in distribution over time since painting (fig. S7).

If the altered flight orientation directions of the migrants with black-painted antennae were a general reflection of the timing of free-running antennal clocks, there should be a correlation between the day of study and the direction of oriented flight; that is, there should be either a progressive increase or a decrease in the angle of orientation with increasing days of study after antennal painting. Because a short free-running period length in DD had been described previously for the timing of adult eclosion behavior in monarchs [figure S3B in (6)] and the *per* and *tim* mRNA patterns measured after 11 days in black-painted antennae were advanced in their peak amounts (Fig. 3C), we calculated the drift in orientation angle over the 14 days of study in a counterclockwise direction relative to mean orientation of control butterflies. Consequently, we observed a significant linear relationship between day of study and orientation angle (Fig. 4B). The slope of the regression line ($y = -9.9x + 279.9$) predicted a free-running period length for the antennal clocks of 23.3 hours, in accordance with the short free-running period found for adult eclosion.

These flight simulator data are consistent with a free-running timing mechanism in the black-painted antennae that influences sun compass orientation. The orientation findings in migrants with black-painted antennae contrast with those of the antennaeless butterflies in which antennal clocks have been removed and no residual group orientation is apparent (Fig. 1B), although these migrants were studied later after antennal removal (fig. S2).

The altered flight directions of migrants with black-painted antennae could also reflect the combined effects of free-running antennal clocks and entrained brain clocks on sun compass orientation. Indeed, flight orientation of migrants with black-painted antennae differed between the LD group and the 6-hour-shifted LD group (Fig. 4A, bottom left and right), suggesting a role of brain clocks. Thus, circadian information from the antennae and the brain may be integrated downstream from the actual clocks themselves, perhaps at an integration site somewhere in the central complex or its output pathways controlling motor behavior.

We found that the antennae are necessary for proper time-compensated sun compass orientation in migratory monarch butterflies. Our results are consistent with a major role of antennal clocks in the timing of Sun compass orientation in migratory monarchs. The antennae may function

alone, without any influence from brain clocks, or antennal output may influence the integration of timing information from brain circadian clocks within the Sun compass structure or at the level of its output pathways. Both possibilities suggest the existence of a crucial but hitherto unknown neural circuit between the antennae and the central complex system.

The role of the antennae in the clock-compass circuitry underlying sun compass orientation could have broad implications, because a similar process may extend to other insects (such as bees, ants, and locusts) that use this orientation mechanism. Furthermore, our results add to the growing list of important nonolfactory functions (e.g., gravity, wind, and sound sensing) housed in the antennae of insects (25, 29, 30).

References and Notes

- L. P. Brower, *J. Lepid. Soc.* **49**, 304 (1995).
- F. A. Urquhart, *The Monarch Butterfly* (Univ. of Toronto Press, Toronto, Canada, 1960).
- O. Froy, A. L. Gotter, A. L. Casselman, S. M. Reppert, *Science* **300**, 1303 (2003).
- H. Mouritsen, B. J. Frost, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 10162 (2002).
- S. M. Perez, O. R. Taylor, R. Jander, *Nature* **387**, 29 (1997).
- H. Zhu *et al.*, *PLoS Biol.* **6**, e4 (2008).
- Q. Yuan, D. Metterville, A. D. Briscoe, S. M. Reppert, *Mol. Biol. Evol.* **24**, 948 (2007).
- H. Zhu *et al.*, *Curr. Biol.* **15**, R953 (2005).
- I. Sauman *et al.*, *Neuron* **46**, 457 (2005).
- J. W. Truman, L. M. Riddiford, *Science* **167**, 1624 (1970).
- P. E. Hardin, *Curr. Biol.* **15**, R714 (2005).
- S. M. Reppert, *Cell* **124**, 233 (2006).
- S. Heinze, U. Homberg, *Science* **315**, 995 (2007).
- M. Sakura, D. Lambrinos, T. Labhart, *J. Neurophysiol.* **99**, 667 (2008).
- G. Liu *et al.*, *Nature* **439**, 551 (2006).
- C. Merlin, M. C. Francois, I. Queguiner, M. Maibeche-Coisne, E. Jacquin-Joly, *Insect Mol. Biol.* **15**, 137 (2006).
- C. Merlin *et al.*, *J. Biol. Rhythms* **22**, 502 (2007).
- J. D. Plautz, M. Kaneko, J. C. Hall, S. A. Kay, *Science* **278**, 1632 (1997).
- J. Schuckel, K. K. Siwicki, M. Stengl, *Cell Tissue Res.* **330**, 271 (2007).
- S. Tanoue, P. Krishnan, B. Krishnan, S. E. Dryer, P. E. Hardin, *Curr. Biol.* **14**, 638 (2004).
- A. S. Saifullah, T. L. Page, *J. Biol. Rhythms* **24**, 144 (2009).
- Materials and methods are available as supporting material on Science Online.
- S. M. Reppert, H. Zhu, R. H. White, *Curr. Biol.* **14**, 155 (2004).
- H. Zhu, R. J. Gegear, A. Casselman, S. Kanginakudru, S. M. Reppert, *BMC Biol.* **7**, 14 (2009).
- S. P. Sane, A. Dieudonne, M. A. Willis, T. L. Daniel, *Science* **315**, 863 (2007).
- J. C. Dunlap, *Cell* **96**, 271 (1999).
- J. Stalleicken *et al.*, *J. Exp. Biol.* **208**, 2399 (2005).
- W. H. Calvert, *J. Lepid. Soc.* **55**, 162 (2001).
- A. Kamikouchi *et al.*, *Nature* **458**, 165 (2009).
- S. Yorozu *et al.*, *Nature* **458**, 201 (2009).
- We thank L. Foley and A. Casselman for assistance; F. Gagnon, C. Collar, and O. Taylor for supplying monarch butterflies; and D. Weaver, Q. Yuan, and members of the Reppert laboratory for their inputs and comments. Supported by NIH grant R01 GM086794.

Supporting Online Material

www.sciencemag.org/cgi/content/full/325/5948/1700/DC1
Materials and Methods
Figs. S1 to S7
References

12 May 2009; accepted 30 June 2009
10.1126/science.1176221